

#### LA-UR-21-20605

Approved for public release; distribution is unlimited.

Title: Sensitivity of modeled fire behavior to small perturbations in initial

conditions

Author(s): Jonko, Alexandra


Yedinak, Kara Conley, Juliana Linn, Rodman Ray Atchley, Adam Lee Parsons, Russell

Intended for: University Seminar at Montana State University, Computer Science

Department, January 25, 2021

Issued: 2021-01-25





# Sensitivity of modeled fire behavior to small perturbations in initial conditions

Alex Jonko<sup>1</sup>, Kara Yedinak<sup>2</sup>, Juliana Conley<sup>1</sup>, Rod Linn<sup>1</sup>, Adam Atchley<sup>1</sup>, and Russ Parsons<sup>3</sup>






<sup>&</sup>lt;sup>1</sup> Computational Earth Science Group, Los Alamos National Laboratory

<sup>&</sup>lt;sup>2</sup> Forest Products Laboratory, US Forest Service

<sup>&</sup>lt;sup>3</sup> Rocky Mountain Research Station, US Forest Service

### Chaotic systems: strong sensitivity to small perturbation

Weather forecasting acknowledges chaotic nature of atmosphere, and uses ensemble simulation




### Chaotic systems: strong sensitivity to small perturbation

European Centre for Medium-Range Weather Forecasts example forecast of temperature anomalies over Europe:


~50 members are used for each forecast

Should we be doing the same for wildfire?




#### The spectrum of fire models

empirical models (e.g. Rothermel)



Fast, simplified, operationally used

"physics-based" models (e.g. FIRETEC)




Detailed, computationally expensive, used in research

## HIGRAD/FIRETEC: coupled fire-atmosphere modeling

#### **HIGRAD/FIRETEC:**

- High GRADient hydrodynamic model of atmospheric flow
- combustion and heat transfer model
- Discrete fuel beds at meter resolution





## HIGRAD/FIRETEC: coupled fire-atmosphere modeling

 Research tool that can be used for simplified simulations of certain aspects of fire behavior, or portions of real wildfires

**Movie removed for RASSTI** 



### HIGRAD/FIRETEC: coupled fire-atmosphere modeling

- Research tool that can be used for simplified simulations of certain aspects of fire behavior, or portions of real wildfires
- Has been used to inform prescribed fire planning
- Internal application: evaluate fire risk to LANL infrastructure


Real Topo Area G, 2 m/s wind coming form west, 4-line ignition. Mown grass around the dome

Movie removed for RASSTI

Are fires chaotic systems?


FIRETEC simulations of the RxCADRE fire experiment, using data from adjacent anemometers, show strong sensitivity of fire to perturbations in the turbulent wind field





#### Are fires chaotic systems?

FIRETEC simulations of the RxCADRE fire experiment, using data from adjacent anemometers, show strong sensitivity of fire to perturbations in the turbulent wind field



Linn, R. et al. (2021) Modeling Low Intensity Fire: Lessons Learned from 2012 RxCADRE. MDPI Atmosphere (in press).

#### **Study Design**

1. Sensitivity to perturbations in atmospheric initial conditions



**Fuels:** homogenous grass

**Winds:** log profile: 2.5 m/s at 10 m

**Vary:** - location of ignition within turbulent

wind field

- fuel moisture

2. Sensitivity to perturbations in initial conditions of fuels




heterogenous Ponderosa Pine stands

log profile: 6.5 m/s at 25 m

- level of aggregation of trees
- location of ignition in heterogeneous fuel beds
- timing of ignition in turbulent wind field

#### 1. Spin up a large turbulent wind field




- Turbulent wind spin up on large (1200 m x 1200 m) domain
- 9 small (400 m x 400 m) domains nested within for fire runs


#### Movie:

- 2 m horizontal wind speed across large domain
- small domain boundaries and ignition lines added for reference


#### 1. Spin up a large turbulent wind field




#### 2. Ignite fires in each of the 9 small domains



- Winds in each small domain do not interact with winds in neighboring domains.
- Fire in one domain is independent of the others.




- Final fire scars from the first 9 simulations highlight variability.
- Next, we run 9 more fires in the same domains, moving ignition forward in time by 60 seconds.
- Since fuels are homogenous, we consider location and timing of ignition as interchangeable.





al fire scars from the first 9 nulations highlight variability.


ext, we run 9 more fires in the ne domains, moving ignition ward in time by 60 seconds.

nce fuels are homogenous, we nsider location and timing of nition as interchangeable.












#### 3. Examine fire outcomes: Clustering and diverging behavior




Impacts of wind at time of ignition




#### The smallest fire in the ensemble is associated with low mean winds in x direction at the inlet




#### The largest fire in the ensemble is associated with high mean winds in x direction at the inlet




Impacts of wind boundary vs initial conditions



Impacts of wind boundary vs initial conditions



Impacts of wind boundary vs initial conditions



Jonko et al. (2021) Sensitivity of grass fires burning in marginal conditions to small perturbations in the turbulent wind field. JGR Atmospheres (submitted).

### Next, let's consider relationships between multiple parameters:



Add in sensitivity to perturbations in fuel initial conditions

#### Heterogeneous fuel beds – Ponderosa Pine




- Tree measurements:
- To populate domains:

Ponderosa Pine stand near Flagstaff, AZ

Simulate 1/f noise process using Inverse Discrete Fourier Transform


#### Levels of fuel aggregation

- Continuous surfaces with power spectral density  $S(f) \sim 1/f^a$
- $a = 0 \rightarrow$  white noise, uncorrelated distribution of trees
- $a = 2 \rightarrow$  Brownian noise, very autocorrelated surface




#### Isolate impacts of turbulence in complex fuels:

Stagger ignitions in time




### **Consider effects** of fuels and **atmosphere:** Low fuel

aggregation




# Consider effects of fuels and atmosphere: Moderate fuel

Moderate fue aggregation



### **Consider effects** of fuels and atmosphere: High fuel

aggregation





#### Overview of some first results

	area burned:	surface fuels consumed:	canopy fuels consumed:
average	19432 m <sup>2</sup> (12%)	3015 kg (6%)	7349 kg (4%)
coefficient of variation	0.43	0.44	0.54
average	22851 m <sup>2</sup> (14%)	3600 kg (8%)	8046 kg (4%)
coefficient of variation	0.35	0.36	0.53
average	32968 m <sup>2</sup> (21%)	5334 kg (11%)	10781 kg (6%)
coefficient of variation	0.40	0.41	0.85

### Back to chaos: transitions between sub-critical and critical states

Example of variable surface to crown fire transition





Example of transition between rapid growth and extinction

